

# Adding Chocolate to Milk Doesn't Take Away Its Nine Essential Nutrients

All milk contains a unique combination of nutrients important for growth and development - including three of the five "nutrients of concern" for which children have inadequate intakes. And, flavored milk accounts for less than 3.5% of added sugar intake in children ages 6-12 and less than 2% in teens.

## 5 Reasons Why Flavored Milk Matters

### 1 KIDS LOVE THE TASTE!

Milk provides nutrients essential for good health and kids will drink more when it's flavored.

### 2 NINE ESSENTIAL NUTRIENTS!

Flavored milk contains the same nine essential nutrients as white milk - calcium, potassium, phosphorous, protein, vitamins A, D and B12, riboflavin and niacin (niacin equivalents) - and is a healthful alternative to soft drinks.

### 3 HELPS KIDS ACHIEVE 3 SERVINGS!

Drinking low-fat or fat-free white or flavored milk helps kids get the 3 daily servings\* of milk recommended by the *Dietary Guidelines for Americans*.

### 4 BETTER DIET QUALITY!

Children who drink flavored milk meet more of their nutrient needs; do not consume more added sugar, fat or calories; and are not heavier than non-milk drinkers.

### 5 TOP CHOICE IN SCHOOLS!

Low-fat chocolate milk is the most popular milk choice in schools and kids drink less milk (and get fewer nutrients) if it's taken away.

These health and nutrition organizations support 3-A-Day of Dairy, a science-based nutrition education program encouraging Americans to consume the recommended three daily servings of nutrient-rich low-fat or fat-free milk and milk products to improve overall health.



[www.nationaldairycouncil.org/childnutrition](http://www.nationaldairycouncil.org/childnutrition)

© National Dairy Council 2009®

#### REFERENCES:

1. NPD Nutrient Intake Database; 2 years ending Feb. 2009.
2. Johnson RK, Frary C, Wang MQ. The nutritional consequences of flavored milk consumption by school-aged children and adolescents in the United States. *J Am Diet Assoc.* 2002;102(6):853-856.
3. National Dairy Council and School Nutrition Association. The School Milk Pilot Test. Beverage Marketing Corporation for National Dairy Council and School Nutrition Association. 2002. [http://www.nutritionexplorations.org/sfs/schoolmilk\\_pilottest.asp](http://www.nutritionexplorations.org/sfs/schoolmilk_pilottest.asp) (Accessed January 4, 2009).
4. NICHD. For Stronger Bones...for Lifelong Health...Milk Matters! Accessed Sept 7, 2009 via [http://www.nichd.nih.gov/publications/pubs/upload/strong\\_bones\\_lifelong\\_health\\_mml.pdf](http://www.nichd.nih.gov/publications/pubs/upload/strong_bones_lifelong_health_mml.pdf)
5. HHS. Best Bones Forever. Accessed Sept 7, 2009 via <http://www.bestbonesforever.gov/>
6. Frary CD, Johnson RK, Wang MQ. Children and adolescents' choices of foods and beverages high in added sugars are associated with intakes of key nutrients and food groups. *J Adolesc Health* 2004;34(1):56-63.
7. American Academy of Pediatrics, Committee on School Health. Soft drinks in schools. *Pediatrics* 2005; 113:152-154.
8. United States Dept. of Health and Human Services, United States Dept. of Agriculture and United States Dietary Guidelines Advisory Committee. *2005 Dietary Guidelines for Americans*. (6th ed. HHS publications, 2005, Washington D.C.)
9. Greer FR, Krebs NF and the Committee on Nutrition. Optimizing bone health and calcium intakes of infants, children and adolescents. *Pediatrics* 2006; 117:578-585.
10. Murphy MM, Douglas JS, Johnson RK, Spence LA. Drinking flavored or plain milk is positively associated with nutrient intake and is not associated with adverse effects on weight status in U.S. children and adolescents. *J Am Diet Assoc* 2008; 108:631-639.
11. Johnson RK, et al. Dietary Sugars Intake and Cardiovascular Health. A Scientific Statement From the American Heart Association. *Circulation*. 2009; 120:1011-1020.
12. ENVIRON International Corporation. School Milk: Fat Content Has Declined Dramatically since the Early 1990s. 2008.
13. Patterson J, Sidel M. The Removal of Flavored Milk in Schools Results in a Reduction in Total Milk Purchases in All Grades, K-12. *J Am Diet Assoc.* 2009; 109(9): A97.

\* DAILY RECOMMENDATIONS - 3 cups of low-fat or fat-free milk or equivalent milk products for those 9 years of age and older and 2 cups of low-fat and fat-free milk or equivalent milk products for children 2-8 years old.